

PEDIATRICS[®]

Discover Pediatric Collections on [COVID-19](#) and [Racism and Its Effects on Pediatric Health](#)

[A statement of reaffirmation for this policy was published at e2021052583](#)

From the American Academy of Pediatrics | Policy Statement

Prevention of Childhood Lead Toxicity

COUNCIL ON ENVIRONMENTAL HEALTH

Pediatrics July 2016, 138 (1) e20161493; DOI: <https://doi.org/10.1542/peds.2016-1493>

[Article](#)

[Figures & Data](#)

[Info & Metrics](#)

[Comments](#)

This article has corrections. Please see:

- COUNCIL ON ENVIRONMENTAL HEALTH. Prevention of Childhood Lead Toxicity. Pediatrics. 2016;38(1):e20161493 - August 01, 2017
- Prevention of Childhood Lead Toxicity - June 01, 2020

 [Download PDF](#)

Abstract

Blood lead concentrations have decreased dramatically in US children over the past 4 decades, but too many children still live in housing with deteriorated lead-based paint and are at risk for lead exposure with resulting lead-associated cognitive impairment and behavioral problems. Evidence continues to accrue that commonly encountered blood lead concentrations, even those below 5 $\mu\text{g}/\text{dL}$ (50 ppb), impair cognition; there is no identified threshold or safe level of lead in blood. From 2007 to 2010, approximately 2.6% of preschool children in the United States had a blood lead concentration $\geq 5 \mu\text{g}/\text{dL}$ (≥ 50 ppb), which represents about 535 000 US children 1 to 5 years of age. Evidence-based guidance is available for managing increased lead exposure in children, and reducing sources of lead in the environment, including lead in housing, soil, water,

and consumer products, has been shown to be cost-beneficial. Primary prevention should be the focus of policy on childhood lead toxicity.

- Abbreviations:

AAP — *American Academy of Pediatrics*

ADHD — *attention-deficit/hyperactivity disorder*

CDC — *Centers for Disease Control and Prevention*

CPSC — *Consumer Product Safety Commission*

EPA — *Environmental Protection Agency*

FDA — *US Food and Drug Administration*

HUD — *Department of Housing and Urban Development*

NHANES — *National Health and Nutrition Examination Survey*

XRF — *x-ray fluorescence*


Overview and Introduction

Primary prevention, reducing or eliminating the myriad sources of lead in the environment of children before exposure occurs, is the most reliable and cost-effective measure to protect children from lead toxicity. Very high blood lead concentrations (eg, >100 µg/dL) can cause significant overt symptoms, such as protracted vomiting and encephalopathy, and even death. Low-level lead exposure, even at blood lead concentrations below 5 µg/dL (50 ppb), is a causal risk factor for diminished intellectual and academic abilities, higher rates of neurobehavioral disorders such as hyperactivity and attention deficits, and lower birth weight in children. No effective treatments ameliorate the permanent developmental effects of lead toxicity. Reducing lead exposure from residential lead hazards, industrial sources, contaminated foods or water, and other consumer products is an effective way to prevent or control childhood lead exposure. Lead poisoning prevention education directed at hand-washing or dust control fails to reduce children's blood lead concentrations. However, pediatricians and parents should be aware of measures to reduce the toxic effects of lead on children, including the promulgation of regulations to screen or test older housing units for lead hazards before occupancy and after major renovation and abatement; revision of federal standards to reduce allowable levels of lead in settled house dust, water, soil, cosmetics, and other consumer products; and enhanced protection for children who live in lead-contaminated communities or near lead-emitting industries.

Scope of the Problem

Over the past 4 decades, blood lead concentrations among US children have declined dramatically since the elimination of lead from gasoline, paints, and other consumer products¹ (Fig 1, Table 1). From 1976 to 1980, blood lead concentrations among US children declined more sharply than anticipated after the phase-out of

leaded gasoline.² In 1978, the US Consumer Product Safety Commission (CPSC) restricted the allowable content of lead in residential paint to 0.06% (600 ppm); in 2008, it was lowered to 0.009% (90 ppm).^{3,4} There have also been significant reductions in tap water lead concentrations since the US Environmental Protection Agency (EPA) promulgated the Lead and Copper Rule.^{5,6} Finally, use of lead solder in canned foods and other consumer products was banned. It is difficult to accurately apportion the decline in blood lead concentrations to specific sources, but the combined effect of these regulations clearly led to the dramatic reductions in children's blood lead concentrations.¹ The key to preventing lead toxicity in children is to reduce or eliminate persistent sources of lead exposure in their environment.



[Download figure](#)

[Open in new tab](#)

[Download powerpoint](#)

FIGURE 1

Timeline of lead poisoning prevention policies and blood lead levels in children aged 1–5 years, by year—NHANES, United States, 1971–2008. BLL, blood lead level; GM BLL, geometric mean blood lead level. Adapted from Brown et al.¹

TABLE 1

Federal Lead Poisoning Prevention Policies

[View inline](#)

Prevention of low-level lead toxicity has historically focused on anticipatory guidance, screening children's blood for lead after exposure, and iron or calcium supplementation to reduce lead absorption.⁷ Unfortunately, studies that evaluated the efficacy of parent education or provision of cleaning equipment to families failed to show significant reductions in children's blood lead concentrations.⁸ Similarly, calcium and iron

supplementation have not consistently been shown to be efficacious in reducing blood lead concentrations of children.^{9,10} Collectively, these studies indicate that the focus of prevention should be on reducing the sources of childhood lead exposures rather than identifying children who have already been unduly exposed or attempting to ameliorate the toxic effects of lead exposure.

In 2005, the American Academy of Pediatrics (AAP) recognized that blood lead concentrations below 10 µg/dL (100 ppb) may impair cognition; no threshold for the toxic effects of lead was identified.⁷ The AAP adopted a blood lead concentration >10 µg/dL (>100 ppb) as the “level of concern” recommended by the Centers for Disease Control and Prevention (CDC), which indicated the need for closer medical and public health management.⁷ Extensive and compelling evidence now indicates that lead-associated cognitive deficits and behavioral problems can occur at blood lead concentrations below 5 µg/dL (50 ppb). In 2012, the US National Toxicology Program of the National Institutes of Health reported that, after other risk factors are accounted for, blood lead concentrations <5 µg/dL (<50 ppb) are strongly associated with intellectual deficits, diminished academic abilities, attention deficits, and problem behaviors (**Table 2**).¹¹ In that same year, the Advisory Committee on Childhood Lead Poisoning Prevention of the CDC concluded that there is no safe level of lead exposure and adopted the use of a reference value of ≥5 µg/dL (≥50 ppb) (based on the 97.5th percentile of blood lead concentrations from the National Health and Nutrition Examination Survey [NHANES]) to be used as a trigger to guide clinical and public health interventions.¹²

TABLE 2

[View inline](#)

Effects of Low-Level Lead Exposure on Academic and Intellectual Abilities, Puberty, Kidney Function, Postnatal Growth, Hearing, and Other Health Endpoints

Low-level elevations in children’s blood lead concentrations, even at concentrations below 5 µg/dL (50 ppb), can result in decrements in cognitive functions, as measured by IQ scores and academic performance.^{13,14} For a given level of exposure, lead-associated IQ decrements are proportionately greater at the lowest blood lead concentrations. The IQ decrement associated with an increase in blood lead concentration from <1 µg/dL (<10 ppb) to 30 µg/dL (300 ppb) was 9.2 IQ points, but the decrement associated with an increase in blood lead concentration from <1 µg/dL (<10 ppb) to 10 µg/dL (100 ppb) was 6.2 IQ points.¹⁴ The population impact of lead on intellectual abilities is substantial. Despite the dramatic reductions in blood lead levels, lead toxicity accounts for an estimated total loss of 23 million IQ points among a 6-year cohort of contemporary US children.¹⁵

Focusing efforts on children who have blood lead concentrations ≥5 µg/dL (≥50 ppb) is efficient but will fail to preserve the majority of lost IQ points in US children. The *prevention paradox* refers to the concept that most disease or disability occurs in low- to moderate-risk groups. Children who have blood lead concentrations ≥5 µg/dL (≥50 ppb) will, on average, experience a lead-associated IQ deficit of 6.1 points, an IQ deficit much larger than that of children who have lower blood lead concentrations (**Fig 2**). Still, if the focus is only on reducing exposures for children who have a blood lead concentration ≥5 µg/dL (≥50 ppb), we will fail to preserve more

than 20 million (>80% of total) of the 23 million IQ points lost among US children with lower lead exposure because there are so many more children who have low to moderate blood lead concentrations (**Fig 2**). No therapeutic interventions currently exist for low blood lead concentrations; therefore, prevention of exposure is paramount. For these reasons, this statement focuses heavily on how pediatricians can help *prevent* lead exposure in children.

[Download figure](#)

[Open in new tab](#)

[Download powerpoint](#)

FIGURE 2

Prevention paradox. The majority of IQ points lost due to lead exposure occur in children who have low to moderate blood lead levels. Using the current reference value of 5 µg/dL, we will protect only 3.1 million IQ points (about 13% of the total). Adapted from Bellinger.¹⁵

Elevated blood lead concentrations can result in the development of behavioral problems in children, including inattention, impulsivity, aggression, and hyperactivity.¹⁶⁻¹⁸ In a nationally representative study of 8- to 15-year-old US children, Froehlich et al¹⁷ found that having a blood lead concentration >1.3 µg/dL (>13 ppb) was associated with an elevated risk for attention-deficit/hyperactivity disorder (ADHD). Children with a blood lead concentration in the lowest tertile (<0.7 µg/dL, or <7 ppb) exhibited, on average, 1 symptom of ADHD,

whereas children with a blood lead concentration in the highest tertile ($>1.3 \mu\text{g/dL}$, or $>13 \text{ ppb}$) exhibited 3 symptoms. Some critics have argued that these “subtle” shifts in behavioral symptoms are inconsequential, but this shift in the population distribution of ADHD symptoms led to an increase in the percentage of children who met criteria for ADHD from 5% to 13%. Approximately 1 in 5 cases of ADHD among US children have been attributed to lead exposure.¹⁷

Antisocial behaviors, including conduct disorder, delinquency, and criminal behaviors, can result from a variety of risk factors, but there is substantial evidence that lead toxicity is 1 of the major risk factors for their development.^{16,19–22} Needleman et al¹⁶ found that adolescents who had higher bone lead concentrations had higher scores for delinquency and aggression. In a meta-analysis of 16 studies, Marcus et al²² concluded that lead exposure, measured via blood lead or bone lead concentrations, was a risk factor for conduct disorder. In 2 prospective longitudinal studies, higher childhood blood lead or tooth lead concentrations resulted in higher rates of self-reported delinquent behaviors and arrests or convictions.^{20,21} Reyes²³ concluded that the reduction in population mean blood lead concentrations was the major risk factor associated with the decline in severe violent behaviors over the past 3 decades.

Limited evidence implicates lead exposure in diminished kidney function in adolescents at low levels of exposure.¹¹ Using the NHANES, Fadrowski et al²⁴ found that, among 769 adolescents with a median blood lead concentration of $1.5 \mu\text{g/dL}$ (15 ppb), a doubling of the concentration led to a significant reduction in the glomerular filtration rate. It is not clear whether chronic, low-level lead exposure in childhood or adolescence is sufficient to result in chronic renal failure or whether it is the cumulative effect of a variety of risk factors that ultimately results in the development of chronic renal failure. Still, this study is consistent with others linking lead exposure with chronic renal failure in adults.¹¹

Lead can cause spontaneous abortion, low birth weight, and reduced growth in children. In a case-control study of pregnant women in Mexico City with blood lead concentrations that ranged from $1.3 \mu\text{g/dL}$ (13 ppb) to $29 \mu\text{g/dL}$ (290 ppb), the odds for spontaneous abortion increased by 1.8 for every $5\text{-}\mu\text{g/dL}$ (50-ppb) increase in maternal blood lead concentration.²⁵ Early studies that examined the association of prenatal lead exposure and low birth weight or preterm birth, measured via either maternal or cord blood lead concentrations, found inconsistent results. However, in a large cohort involving more than 34 000 live births, investigators found that a $5\text{-}\mu\text{g/dL}$ (50-ppb) increase in blood lead concentrations was associated with a 61-g decrement in birth weight.²⁶ The National Toxicology Program concluded that maternal blood lead concentrations $<5 \mu\text{g/dL}$ ($<50 \text{ ppb}$) are associated with lower birth weight.

Preventing Lead Toxicity

Despite historical reductions in children’s blood lead concentrations, preventing childhood lead toxicity remains a major public health priority in the United States. Many children who live in older, poorly maintained

housing or older housing that undergoes renovation are at high risk for lead exposure. In the NHANES conducted from 2007 to 2010, approximately 2.6% of preschool children in the United States had a blood lead concentration ≥ 5 $\mu\text{g}/\text{dL}$ (≥ 50 ppb), which represents about 535 000 US children 1 to 5 years of age.¹² Children who lived in older housing units experienced an increased risk for having a blood lead concentration in excess of 5 $\mu\text{g}/\text{dL}$ (50 ppb); 15% of US children who lived in housing units built before 1950 had a blood lead concentration ≥ 5 $\mu\text{g}/\text{dL}$ (≥ 50 ppb), whereas 4.2% of children who lived in housing built between 1950 and 1978 had a blood lead concentration ≥ 5 $\mu\text{g}/\text{dL}$ (≥ 50 ppb), compared with 2.1% of children who lived in housing units built after 1978.²⁷ No treatments have been shown to be effective in ameliorating the permanent developmental effects of lead toxicity.²⁸ Finally, the economic costs of childhood lead toxicity are substantial. Despite the historical reductions in blood lead concentrations, it has been estimated that the annual cost of childhood lead exposure in the United States is \$50 billion.²⁹ For every \$1 invested to reduce lead hazards in housing units, society would benefit by an estimated \$17 to \$221, a cost–benefit ratio that is comparable with the cost–benefit ratio for childhood vaccines.³⁰

The key to preventing lead toxicity in children is identification and elimination of the major sources of lead exposure. Primary prevention of lead exposure is now widely recognized as the optimal strategy because of the irreversible effects of low-level lead toxicity.^{7,12} The primary prevention approach contrasts with practices and policies that too often have relied predominantly on detection of lead exposure only after children develop elevated blood lead concentrations.

Sources and Variability of Lead Exposure

Lead ingestion and absorption are dynamic during the first 2 years of life. Blood lead concentrations of children who live in lead-contaminated environments typically increase rapidly between 6 and 12 months of age, peak between 18 and 36 months of age, and then gradually decrease.³¹ The peak in children's blood lead concentrations stems from the confluence of normal mouthing behaviors and increasing mobility.³¹ Younger children also absorb lead more efficiently than older children and adults.³² Iron deficiency can also increase the absorption of lead.³³

A large number of housing units in the United States contain lead-based paint. In a national survey of housing conducted in 2011, it was estimated that 37 million (35%) of 106 million housing units contain lead-based paint.³⁴ Lead-based paint is the most common, highly concentrated source of lead exposure for children who live in older housing.³⁵ Paint that was used on both the interior and exterior of houses through the 1950s contained higher concentrations of lead than that of houses built in later years.^{34,35} The lead concentration in paint and other media can be measured by using a hand-held instrument called the x-ray fluorescence (XRF) spectrum analyzer or by chemically analyzing paint chips.

The US Department of Housing and Urban Development (HUD) defines lead-based paint as an XRF reading ≥ 1 $\mu\text{g}/\text{cm}^2$ or 5000 ppm of lead in a paint chip.³⁶ The presence of *lead-based paint* is not as predictive of childhood lead exposure as a *lead paint hazard*. A lead paint hazard is defined by the EPA as “any condition that causes exposure to lead from contaminated dust, lead-contaminated soil, or lead-contaminated paint that is deteriorated, or the presence of accessible (or chewable) surfaces, friction surfaces or impact surfaces that would result in adverse human health effects.”³⁷

Age of the housing is a major determinant of lead paint hazards. For housing built from 1978 to 1998, 2.7% contained one or more lead paint hazards, whereas the prevalence of residential hazards increased to 11.4% of housing built from 1960 to 1977, 39% of housing built from 1940 to 1959, and 67% of housing units built before 1940.³⁴ Federal regulations for defining a lead paint hazard in house dust are obsolete. Federal agencies have set environmental lead standards to protect children from having a blood lead concentration ≥ 10 $\mu\text{g}/\text{dL}$ (≥ 100 ppb), but it is now recognized that there is no safe level of lead exposure. Therefore, because the current standards for lead in house dust, water, and soil remain too high to protect children,^{31,38} the percentage of housing that contains one or more lead paint hazards described above is an underestimate.

Lead-based paint is the major source of lead, but ingestions of lead-contaminated house dust and residential soil are the major pathways for exposure (**Fig 3**).³⁵⁻⁴² House dust, which can be contaminated by small particles of lead-based paint or track-in of lead-contaminated soil, is a major pathway of lead exposure for children who live in older, poorly maintained housing.⁴⁰ Ingestions of lead-contaminated house dust and soil are also the primary pathways of exposure for children who live in homes that were recently abated or renovated.⁴³⁻⁴⁵

[Download figure](#)

[Open in new tab](#)

[Download powerpoint](#)

FIGURE 3

Contribution of lead exposure to children's blood lead concentrations. Adapted from Lanphear et al.³¹ and Spanier et al.⁴⁵

Sampling house dust for lead hazards involves using a special wipe to sample a specified area, such as the floor, which is readily accessible to a child, or a window sill or window trough.³⁶ Windows are often more heavily contaminated than floors because exterior paints often contained higher concentrations of lead, and window troughs can act as reservoirs. Sampling house dust for lead is used to screen older housing units that may contain lead hazards at the time of purchase or rental and before occupancy; to conduct a full risk assessment that involves extensive sampling of settled dust in housing units that failed a lead hazard screen or where there is a high probability of a lead hazard; and to conduct clearance testing after repair or renovation of painted surfaces and after lead abatement, to verify that the housing unit is safe for occupancy (Table 3).³⁸

TABLE 3

Common Sources of Lead Exposure

[View inline](#)

Lead-contaminated soil is an important source of lead intake for children.^{40,41} Lead-contaminated soil can directly contribute to children's blood lead concentrations via soil ingestion and indirectly from soil tracked indoors on shoes, which then contaminates house dust (**Fig 3**). Former mine and smelter communities present a particular risk to children for the ingestion of lead-contaminated soil, but lead in urban soil also is often heavily contaminated from the past use of leaded gasoline and paints. Other sources of lead in soil include weathering of lead-based exterior paint and nearby renovation or demolition activity. Soil testing is usually performed in areas where children play and the foundation perimeter. The EPA standards are 400 µg of lead per gram of soil for play areas and 1200 µg/g for the foundation perimeter.³⁷ Children's blood lead concentrations increase by approximately 3.8 µg/dL (38 ppb) for every 1000-ppm increase in soil lead concentration.⁴⁰

Water is an important but often overlooked source of exposure for children, especially for infants who are formula fed.^{5,46,47} Water typically contributes to approximately 20% of a child's blood lead concentrations if the water lead concentration exceeds 5 ppb (**Fig 3**).³¹ The contribution of lead from water can be much higher for some children, especially for infants who ingest large quantities of tap water.^{5,46,47} Children who reside in communities with lead service lines and inadequate anticorrosion control are also at increased risk for elevated blood lead concentrations.⁴⁸

Phasing out leaded gasoline and creating stricter national air lead standards led to large reductions in the contribution of airborne lead to children's blood lead concentrations. Still, in some communities, such as those surrounding regional airports, airborne lead is an important source of lead exposure. Airborne lead is ingested primarily after it settles in house dust and soil where children play. Current sources of airborne lead include lead battery recycling operations, piston engine aircraft, and incinerators.⁴⁹ The contributions of airborne lead to children's blood lead concentrations are proportionately greater at the lower levels of exposure than at higher levels.⁴⁹

Other sources of lead intake for children have been identified, such as nutritional supplements and folk medicines, ceramic dishware, and cosmetics⁵⁰⁻⁵² (**Table 3**). Lead brought into the home from a worksite by a parent can also be a major source of exposure for some children.⁵³ Consumer products such as children's toys, lunch boxes, crayons, and lipstick that are contaminated with lead have received a great deal of attention. These products constitute a small source of lead intake for most children, but they can be the major source for an individual child. Moreover, because lead exposure is cumulative and there is no apparent threshold for the adverse effects of lead exposure, all sources of lead exposure should be eliminated. It is the responsibility of the relevant federal agencies, such as the CPSC and the Food and Drug Administration (FDA), to promulgate and enforce standards that will protect children from lead-contaminated consumer products.

Residential Standards for Lead in Paint, Dust, and Water

Lead in Paint and Dust

Under section 403 of Title X, the US Congress mandated the EPA to promulgate residential health-based lead standards that are designed to protect children from lead toxicity.³⁷ Standards are necessary to identify lead hazards before a child is unduly exposed and to identify the source of lead exposure for children who have blood lead concentrations ≥ 5 $\mu\text{g}/\text{dL}$ (≥ 50 ppb).³¹ Unless performed carefully, attempts to reduce lead exposure, such as abatement, repair, or renovation, can result in increased contamination and elevation in a child's blood lead concentration.⁴³⁻⁴⁵ Dust clearance tests, which involve collecting dust from floors or windows of a home by using a lead-free material that resembles a baby wipe, should be conducted after extensive repair, renovation, or abatement of older housing units to determine whether the housing intervention was sufficient to protect children from lead hazards, especially in housing units built before 1960.^{27,34} Property owners are required to disclose possible presence of lead-based paint in properties built before 1978 and are required to provide the blue pamphlet from the EPA, HUD, and Consumer Product Safety Commission titled "Protect Your Family From Lead in Your Home" at the time of rental or sale.

Most existing lead standards fail to protect children (**Table 4**). In 1978, the CPSC set the maximum paint lead concentration at 0.06% (600 ppm), because there was evidence that paint could be manufactured with this lower level of contamination.³ Similarly, the EPA's action level of 15 ppb of lead in water, which is used to regulate water systems in the United States, is routinely (but erroneously) used as a health-based standard; it was not intended as a health-based standard, nor does it adequately protect children or pregnant women from adverse effects of lead exposure.^{5,31} In 1988, the HUD established a postabatement floor dust standard of 200 $\mu\text{g}/\text{ft}^2$ because there was evidence that it was feasible to attain, not because it was demonstrated to be safe or protective. In 2001, the EPA promulgated residential lead standards of 40 $\mu\text{g}/\text{ft}^2$ for floors and 250 $\mu\text{g}/\text{ft}^2$ for window sills.³⁷ Unfortunately, these standards, which failed to protect children from having a blood lead concentration ≥ 10 $\mu\text{g}/\text{dL}$ (≥ 100 ppb) when they were first promulgated, dictate the levels of lead contamination considered "normal" or "low," and they provide an illusion of safety.^{38,40} At a floor standard of 40 $\mu\text{g}/\text{ft}^2$, the current EPA standard for floors, 50% of children were estimated to have a blood lead concentration ≥ 5 $\mu\text{g}/\text{dL}$ (≥ 50 ppb); 5% of children have a blood lead concentration ≥ 5 $\mu\text{g}/\text{dL}$ (≥ 50 ppb) at a median floor dust lead level of 1.5 $\mu\text{g}/\text{ft}^2$ (**Fig 4**).⁴²

TABLE 4

Federal Standards for Lead in House Paint, House Dust, Soil, Water, Air, and Candy

[View inline](#)

[Download figure](#)

[Open in new tab](#)

[Download powerpoint](#)

FIGURE 4

Estimated probability of blood lead concentrations ≥ 5 $\mu\text{g}/\text{dL}$ for children living in pre-1978 housing by floor dust lead level, NHANES, 1999–2004. CI, confidence interval. Adapted from Dixon et al.⁴²

Scraping, sanding, or construction during painting, repair, renovation, or abatement of older housing can result in lead contamination of a child's environment.^{41,43–45,54} In a controlled study of children with baseline blood lead concentrations < 22 $\mu\text{g}/\text{dL}$ (< 220 ppb), Aschengrau et al⁴¹ reported a 6.5- $\mu\text{g}/\text{dL}$ (65-ppb) *increase* in blood lead concentrations for children whose homes had undergone paint abatement. Clark et al⁴⁴ reported that 6-month-old infants were 11 times more likely to have a ≥ 5 $\mu\text{g}/\text{dL}$ (≥ 50 ppb) increase in blood lead concentrations after abatement compared with older children. Spanier et al⁴⁵ reported that routine renovation of older housing was associated with a 12% higher mean blood lead concentration. These studies indicate that the levels of lead-contaminated dust generated by lead hazard control work or housing renovations can result in excessive lead exposure and absorption for children unless there is sufficient cleanup and clearance testing after the work is completed. The HUD has published technical guidelines and regulations for workers involved in lead-based paint abatement or remediation of housing.³⁶

In 1992, the US Congress mandated the EPA to promulgate regulations to protect children from lead exposure resulting from housing repairs and renovation.³⁷ In 2011, the EPA finalized recommendations for the Lead Renovation, Repair and Painting Rule.⁵⁴ Unfortunately, the EPA failed to recommend the validated wipe-sampling method for clearance testing. Instead, it used an unvalidated cloth test, which should not be confused with the validated wipe sampling test. The white cloth test assumes that if dust is visible on a white cloth (ie, the “white glove test”), it contains a lead hazard; conversely, if there is no visible dust, it does not contain a lead hazard.⁵⁴ Although it would be valuable to have a quick test to identify the presence of a lead hazard, the white cloth test is not a validated tool and is not a reliable way to quantify the presence of a lead hazard.

Lead hazard control work can result in sizable reductions in the magnitude of dust lead loading when proper procedures are followed and cleanup and postwork clearance testing are performed. In 1 study, dust lead levels (measured as micrograms of lead per area) immediately after professional abatement were 8.5 $\mu\text{g}/\text{ft}^2$, 8.0 $\mu\text{g}/\text{ft}^2$, and 21 $\mu\text{g}/\text{ft}^2$ for floors, interior window sills, and window troughs, respectively, representing reductions of more than 80% compared with preabatement levels.⁵⁵ In another study of more than 2600 housing units, postabatement dust lead levels were 12 $\mu\text{g}/\text{ft}^2$, 31 $\mu\text{g}/\text{ft}^2$, and 32 $\mu\text{g}/\text{ft}^2$ for floors, window sills and window troughs, respectively.⁵⁶ These levels were achieved with dust clearance testing set at 100 $\mu\text{g}/\text{ft}^2$ or higher, but floor dust lead levels below 5 $\mu\text{g}/\text{ft}^2$ can be achieved by following a specific protocol. In 1 unpublished study of more than 160 housing units built before 1978, 1 group found that it is possible to routinely meet floor lead levels below 5 $\mu\text{g}/\text{ft}^2$ after housing renovations costing an average of \$5600 (B. Lanphear, MD, MPH, Simon Fraser University, unpublished data).

Lead in Water

The primary sources of lead in water, which can be dissolved or particulate, consist of lead service lines, lead solder, and brass fittings that contain high concentrations of lead.⁵ Plumbing installed before 1986, the year a federal ban was issued on using lead pipe and lead solder and a maximum lead content of 8% by weight for brass plumbing was established, is more likely to contain higher concentrations of lead.⁵ Lead services lines that are being replaced, are undergoing maintenance, or are damaged can release particles of lead that can be ingested.⁵⁷ Partial service line replacement, which is sometimes performed to minimize the cost of service line repair by water authorities, fails to reduce lead exposure.⁵⁷ Proper maintenance and ultimately full replacement of water service lines will be necessary to eliminate lead intake from water, but it must be performed with proper precautions. In the interim, water filters that are certified by the National Sanitation Foundation for lead removal can effectively reduce water lead concentrations. The EPA recommends running the cold water of residential units for up to 2 minutes to flush the lead leached from pipes out of the plumbing system, but flushing is useful only in housing units without lead service lines.⁵⁸⁻⁶¹ In housing units without lead service lines, and where the primary source is brass fittings or lead-soldered joints, a 1-minute flush may be sufficient, depending on the length of plumbing; for housing units with lead service lines, flushing may *increase* lead exposure, again depending on the length of the lead service lines.⁵⁸⁻⁶¹

Drinking fountains in older schools can be an important source of lead exposure.⁵ Unfortunately, there are no regulations for evaluating lead contamination of school drinking fountains in most states.

Implementation of the Lead and Copper Rule has significantly reduced tap water lead levels. In 1991, the US EPA set an action level for lead in water of 15 µg/L or (15 ppb).⁶ Communities in which >10% of water samples taken from various taps throughout the system exceed 15 ppb are considered to be out of compliance and are required by the EPA to take action to reduce lead levels using corrosion control methods or replacement of lead service lines. The action level is used as an administrative tool to evaluate community-level exposure; it is not a health-based standard. The maximum contaminant level goal, the value the EPA deems acceptable for health, is 0.

Testing Asymptomatic Children for Elevated Blood Lead Concentrations

In the primary care office, primary prevention begins with education and counseling. Ideally, environmental assessments, such as screening older housing units, occurs before a child is born so that parents can identify and hire trained workers to abate environmental lead exposure hazards.¹² It is especially important to conduct an environmental assessment for lead if a family resides in a housing unit built before 1960 that has undergone recent renovation, repair, or painting or if it is poorly maintained.

Screening questionnaires frequently used in the primary care setting fail to identify children who have elevated blood lead concentrations,⁶² but they may be useful as a tool to identify lead hazards in children who have a blood lead concentration ≥ 5 µg/dL (≥ 50 ppb). In addition, public health agencies often use other methods of targeting children who should be screened with a blood lead test on the basis of community and residential characteristics, such as older housing. Blood lead surveillance data can be used to identify cities, communities, or housing units at higher than typical risk for lead poisoning. Technologies using geographic information system-based analyses and surveillance from electronic medical records are important tools to identify at-risk children who should have their blood lead concentration measured.

In 1991, the CDC recommended universal blood lead testing for all children.⁶³ In 2005, the AAP recommended that states and cities formulate their own lead screening recommendations on the basis of local data because of the wide variation in lead exposure.⁷ The AAP, consistent with the CDC, recommended universal screening of children's blood for lead if they lived in communities with more than 27% of housing built before 1950 or a prevalence of blood lead concentrations ≥ 10 µg/dL in children 12 to 36 months old of 12% or greater.^{7,12,63,64} Screening is not efficient after 36 months of age unless specific high-risk factors are identified; the likelihood of a child having a blood lead concentration >10 µg/dL after 36 months of age is low.⁶⁵ These recommendations now need to be updated to conform to with our new understanding of lead toxicity.^{11,12}

A detailed evaluation and follow-up of children who have blood lead concentrations <10 µg/dL (<100 ppb) is now indicated. Current federal regulations for clinical laboratory testing through the Clinical Laboratory Improvement Amendments of 1988⁶⁶ permit an allowable laboratory error in blood lead proficiency testing

programs of $\pm 4 \mu\text{g}/\text{dL}$ ($\pm 40 \text{ ppb}$) for blood lead concentrations $\leq 20 \mu\text{g}/\text{dL}$ ($\leq 200 \text{ ppb}$). This range of error can result in children being misclassified and cause additional anxiety or false comfort when blood lead concentrations within the margin of error erroneously are interpreted as going up or down. The majority of laboratories analyzing blood lead reference materials routinely achieved laboratory error of $\pm 2 \mu\text{g}/\text{dL}$ ($\pm 20 \text{ ppb}$) at blood lead concentrations $\leq 20 \mu\text{g}/\text{dL}$ ($\leq 200 \text{ ppb}$).⁶⁷ Changing the allowable laboratory error to tighter performance requirements, such as $\pm 2 \mu\text{g}/\text{dL}$ ($\pm 20 \text{ ppb}$), could decrease misclassification of children and lead to better allocation of health care resources.

Case Management of Children With a Blood Lead Concentration at or Above Reference Value

The AAP is adopting the current reference value of $\geq 5 \mu\text{g}/\text{dL}$ ($\geq 50 \text{ ppb}$) for case management.¹² The CDC recommended that the 97.5th percentile of blood lead concentrations derived from the combination of the 2 most recent cycles of NHANES data be used to identify children who have unacceptably high exposure and to set public health goals.¹² The CDC will reconsider the reference value for children's blood lead concentrations every 4 years.¹²

After confirmatory testing, it is important to monitor children who have blood lead concentrations $\geq 5 \mu\text{g}/\text{dL}$ ($\geq 50 \text{ ppb}$). The pediatrician should inform the local or state health department and request an inspection of the child's house to identify and remediate any lead hazards (**Table 4**). Screening children for iron deficiency and insufficient dietary calcium intake is also important.⁷ A detailed description of the diagnosis and treatment of significant lead toxicity (ie, $\geq 45 \mu\text{g}/\text{dL}$ [$\geq 450 \text{ ppb}$]) is beyond the scope of this policy statement, but guidance is available in an earlier publication of the AAP⁷ and through the Pediatric Environmental Health Specialty Units Web site (www.pehsu.net) (**Table 5**). Children who have elevated blood lead concentrations need to be monitored until environmental investigations and remediation are complete and blood lead concentrations decline.¹²

TABLE 5

[View inline](#)

AAP Recommendations on Management of Childhood Lead Exposure and Poisoning

The AAP recognizes that environmental investigations will typically be conducted by local or state health or environmental departments to identify sources of lead exposure for a child who has a blood lead concentration $\geq 5 \mu\text{g}/\text{dL}$ ($\geq 50 \text{ ppb}$). In many cases, however, the pediatrician can provide clues about possible sources of lead intake by taking a careful history.

Case management involves a thorough investigation of potential sources of lead poisoning in a child's environment, including paint, house dust, water, and soil. Case management also includes a questionnaire and visual inspection for other potential sources of lead exposure, including antique furniture, toys, ethnic folk remedies, and consumer products such as imported food, cosmetics, and ceramics.^{12,50–52} It can include testing deteriorated paint on furniture, such as a crib, taking dust samples from child care settings or a family member's house, and taking soil samples from a child's play area.

Summary and Recommendations

Lead toxicity results in substantial, population-level effects on children's intellectual abilities, academic abilities, problem behaviors, and birth weight. Pediatricians may be well equipped to advocate for more stringent regulations to reduce sources of lead exposure and prevent childhood lead exposure. The AAP recognizes the importance of a variety of educational, enforcement, and environmental actions to reduce the number of children who are exposed to lead hazards and concur with recent detailed recommendations for prioritization of primary prevention of lead toxicity.^{7,12,68-70} The AAP offers the following recommendations for government as well as pediatricians, other health care providers, and public health officials.

Recommendations for Government

1. The federal government should expand the resources currently offered by the HUD to local and state governments for lead hazard control work.
2. The federal government should provide both financial and nonfinancial resources and technical guidance through the CDC, the EPA, and the HUD to state and local public health agencies as well as environmental and housing agencies engaged in childhood lead poisoning prevention efforts.
3. The US EPA and HUD should review their protocols for identifying and mitigating residential lead hazards (eg, lead-based paint, dust, and soil) and lead-contaminated water from lead service lines or lead solder and revise downward the allowable levels of lead in house dust, soil, paint, and water to conform with the recognition that there are no safe levels of lead.
4. The federal government should resume and expand its vital role in providing federal public health leadership in childhood lead poisoning prevention work through the CDC. Allocation of additional resources would be necessary to accomplish this goal.
5. The Centers for Medicare & Medicaid Services, which is responsible for regulating clinical laboratory testing through the Clinical Laboratory Improvement Amendments of 1988,⁶⁹ should expeditiously revise current regulations for allowable laboratory error permitted in blood lead proficiency testing programs from ± 4 $\mu\text{g}/\text{dL}$ (± 40 ppb) to ± 2 $\mu\text{g}/\text{dL}$ (± 20 ppb) for blood lead concentrations ≤ 20 $\mu\text{g}/\text{dL}$ (≤ 200 ppb).¹² In the future, when feasible, allowable laboratory error permitted in blood lead proficiency testing programs should be reduced even more, to ± 1 $\mu\text{g}/\text{dL}$ (± 10 ppb) for blood lead concentrations ≤ 20 $\mu\text{g}/\text{dL}$ (≤ 200 ppb).
6. The federal government should continue to conduct the NHANES and provide national data on trends in blood lead concentrations. These newer data should be used by the CDC to periodically formulate a new reference value and guide clinical and public health interventions.
7. The federal government should continue to regularly survey children and adolescents in the NHANES for ADHD and conduct disorder by using validated diagnostic surveys from the *Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition* to examine the association of lower blood lead concentrations with these conditions.
8. Local or state governments, in consultation with pediatricians, should develop policies and regulations requiring the remediation of lead-contaminated housing and child care facilities, including the elimination of

lead hazards during transfer of rental units or renovation or demolition of older housing.

9. State and local governments should collect, analyze, and publish blood lead test results performed in their jurisdictions and should regularly publish reports of age of housing and other risk factors for children having blood lead concentrations ≥ 5 $\mu\text{g}/\text{dL}$ (≥ 50 ppb). These reports should be readily available to pediatricians, health care providers, and the public.
10. Federal, state, and local governments should provide resources for environmental evaluations and case management of children who have blood lead concentrations ≥ 5 $\mu\text{g}/\text{dL}$ (≥ 50 ppb), in conjunction with the child's primary care provider.
11. State and local governments should take steps to ensure that water fountains in schools do not exceed water lead concentrations of 1 ppb.

Recommendations for Pediatricians, Health Care Providers, and Public Health Officials

1. Pediatricians are in a unique position to work with public health officials to conduct surveys of blood lead concentrations among a randomly selected, representative sample of children in their states or communities at regular intervals to identify trends in blood lead concentrations. These periodic surveys are especially important for children who live in highly contaminated communities, such as smelter communities or regions with a historically high prevalence of lead exposure.
2. Pediatricians, health care providers, and public health officials should routinely recommend individual environmental assessments of older housing,¹² particularly if a family resides in a housing unit built before 1960 that has undergone recent renovation, repair, or painting or that has been poorly maintained.
3. Pediatricians and public health officials should advocate for the promulgation and enforcement of strict legal standards based on empirical data that regulate allowable levels of lead in air, water, soil, house dust, and consumer products. These standards should address the major sources of lead exposure, including industrial emissions, lead paint in older housing, lead-contaminated soil, water service lines, and consumer products.
4. Pediatricians should be familiar with collection and interpretation of reports of lead hazards found in house dust, soil, paint, and water, or they should be able to refer families to a pediatrician, health care provider, or specialist who is familiar with these tools.
5. Pediatricians, women's health care providers, and public health officials should be familiar with federal, state, local, and professional recommendations or requirements for screening children and pregnant women for lead poisoning.^{12,68,69}
6. Pediatricians and other primary care providers should test asymptomatic children for elevated blood lead concentrations according to federal, local, and state requirements. Immigrant, refugee, and internationally adopted children also should be tested for blood lead concentrations when they arrive in the United States because of their increased risk.^{71,72} Blood lead tests do not need to be duplicated, but the pediatrician or other primary care provider should attempt to verify that screening was performed elsewhere and determine the result before testing is deferred during the office visit.
7. Pediatricians and other primary care health providers should conduct targeted screening of children for elevated blood lead concentrations if they are 12 to 24 months of age and live in communities or census

block groups with $\geq 25\%$ of housing built before 1960 or a prevalence of children's blood lead concentrations $\geq 5 \mu\text{g/dL}$ ($\geq 50 \text{ ppb}$) of $\geq 5\%$.

8. Pediatricians and other primary care providers should test children for elevated blood lead concentrations if they live in or visit a home or child care facility with an identified lead hazard or a home built before 1960 that is in poor repair or was renovated in the past 6 months.^{7,12}
 9. Pediatricians and primary care providers should work with their federal, state, and local governments to ensure that a comprehensive environmental inspection is conducted in the housing units of children who have blood lead concentrations $\geq 5 \mu\text{g/dL}$ ($\geq 50 \text{ ppb}$) and that they receive appropriate case management.
-

LEAD AUTHOR

Bruce Perrin Lanphear, MD, MPH, FAAP

COUNCIL ON ENVIRONMENTAL HEALTH EXECUTIVE COMMITTEE, 2015–2016

Jennifer A. Lowry, MD, FAAP, Chairperson

Samantha Ahdoot, MD, FAAP

Carl R. Baum, MD, FACMT, FAAP

Aaron S. Bernstein, MD, MPH, FAAP

Aparna Bole, MD, FAAP

Heather Lynn Brumberg, MD, MPH, FAAP

Carla C. Campbell, MD, MS, FAAP

Bruce Perrin Lanphear, MD, MPH, FAAP

Susan E. Pacheco, MD, FAAP

Adam J. Spanier, MD, PhD, MPH, FAAP

Leonardo Trasande, MD, MPP, FAAP

FORMER EXECUTIVE COMMITTEE MEMBERS

Kevin C. Osterhoudt, MD, MSCE, FAAP

Jerome A. Paulson, MD, FAAP

Megan T. Sandel, MD, MPH, FAAP

CONTRIBUTOR

Paul Thomas Rogers, MD, FAAP

LIAISONS

John M. Balbus, MD, MPH – *National Institute of Environmental Health Sciences*

Todd A. Brubaker, DO – *Section on Medical Students, Residents, and Fellowship Trainees*

Nathaniel G. DeNicola, MD, MSc – *American College of Obstetricians and Gynecologists*

Ruth Ann Etzel, MD, PhD, FAAP – *US Environmental Protection Agency*

Mary Ellen Mortensen, MD, MS – *CDC/National Center for Environmental Health*

Mary H. Ward, PhD – *National Cancer Institute*

STAFF

Paul Spire

Footnotes

- Accepted May 5, 2016.
- This document is copyrighted and is property of the American Academy of Pediatrics and its Board of Directors. All authors have filed conflict of interest statements with the American Academy of Pediatrics. Any conflicts have been resolved through a process approved by the Board of Directors. The American Academy of Pediatrics has neither solicited nor accepted any commercial involvement in the development of the content of this publication.
- Policy statements from the American Academy of Pediatrics benefit from expertise and resources of liaisons and internal (AAP) and external reviewers. However, policy statements from the American Academy of Pediatrics may not reflect the views of the liaisons or the organizations or government agencies that they represent.
- The guidance in this statement does not indicate an exclusive course of treatment or serve as a standard of medical care. Variations, taking into account individual circumstances, may be appropriate.

- All policy statements from the American Academy of Pediatrics automatically expire 5 years after publication unless reaffirmed, revised, or retired at or before that time.
- **FINANCIAL DISCLOSURE:** The authors have indicated they have no financial relationships relevant to this article to disclose.
- **FUNDING:** No external funding.
- **POTENTIAL CONFLICT OF INTEREST:** The authors have indicated they have no potential conflicts of interest to disclose.

References

1. ↗Brown MJ, Margolis S; Centers for Disease Control and Prevention. Lead in drinking water and human blood lead levels in the United States. *MMWR Suppl*. 2012;**61**(4 suppl 1):1–9pmid:22874873 [PubMed](#) [Google Scholar](#)
2. ↗Annest JL, Pirkle JL, Makuc D, Neese JW, Bayse DD, Kovar MG. Chronological trend in blood lead levels between 1976 and 1980. *N Engl J Med*. 1983;**308**(23):1373–1377pmid:6188954 [CrossRef](#) [PubMed](#) [Google Scholar](#)
3. ↗Committee on Toxicology, Assembly of Life Sciences, National Research Council. Recommendations for the prevention of lead poisoning in children. *Nutr Rev*. 1976;**34**(11):321–327pmid:1004799 [Abstract/FREE Full Text](#) [Google Scholar](#)
4. ↗Consumer Product Safety Commission. Final rule. Children's products containing lead; determinations regarding lead content limits on certain materials or products. *Fed Regist*. 2009;**74**(164):43031–43042 Available at: <https://www.cpsc.gov/PageFiles/77828/leadcontent.txt>. Accessed January 14, 2016 [Google Scholar](#)
5. ↗Triantafyllidou S, Edwards M. Lead (Pb) in tap water and in blood: implications for lead exposure in the United States. *Crit Rev Environ Sci Technol*. 2012;**42**(13):1297–1352 [CrossRef](#) [Google Scholar](#)
6. ↗US Environmental Protection Agency. Drinking water regulations: maximum contaminant level goals and national primary drinking water regulations for lead and copper; Final Rule. *Fed Regist*. 1991;**56**(11):26460–26564 [Google Scholar](#)
7. ↗American Academy of Pediatrics Committee on Environmental Health. Lead exposure in children: prevention, detection, and management. *Pediatrics*. 2005;**116**(4):1036–1046pmid:16199720 [Abstract/FREE Full Text](#) [Google Scholar](#)
8. ↗Yeoh B, Woolfenden S, Lanphear B, Ridley GF, Livingstone N. Household interventions for preventing domestic lead exposure in children. *Cochrane Database Syst Rev*. 2012;**4**(4):CD006047pmid:22513934 [PubMed](#) [Google Scholar](#)
9. ↗Rico JA, Kordas K, López P, et al. Efficacy of iron and/or zinc supplementation on cognitive performance of lead-exposed Mexican schoolchildren: a randomized, placebo-controlled trial. *Pediatrics*. 2006;**117**(3). Available at: www.pediatrics.org/cgi/content/full/117/3/e518pmid:16510631 [Abstract/FREE Full Text](#) [Google Scholar](#)
10. ↗Sargent JD, Dalton MA, O'Connor GT, Olmstead EM, Klein RZ. Randomized trial of calcium glycerophosphate-supplemented infant formula to prevent lead absorption. *Am J Clin Nutr*. 1999;**69**(6):1224–1230pmid:10357743 [Abstract/FREE Full Text](#) [Google Scholar](#)
11. ↗National Toxicology Program. *Monograph on Health Effects of Low-Level Lead*. Research Triangle Park, NC: National Institute of Environmental Health Sciences; 2012:xiii, xv–148 [Google Scholar](#)

12. [↗](#)Centers for Disease Control and Prevention, Advisory Committee on Childhood Lead Poisoning Prevention. *Low Level Lead Exposure Harms Children: A Renewed Call for Primary Prevention*. Atlanta, GA: Centers for Disease Control and Prevention; 2012. Available at: www.cdc.gov/nceh/lead/ACCLPP/Final_Document_030712.pdf. Accessed January 14, 2016 [Google Scholar](#)
13. [↗](#)Lanphear BP, Dietrich K, Auinger P, Cox C. Cognitive deficits associated with blood lead concentrations <10 microg/dL in US children and adolescents. *Public Health Rep*. 2000;**115**(6):521–529pmid:11354334 [CrossRef](#) [PubMed](#) [Google Scholar](#)
14. [↗](#)Lanphear BP, Hornung R, Khoury J, et al. Low-level environmental lead exposure and children's intellectual function: an international pooled analysis. *Environ Health Perspect*. 2005;**113**(7):894–899pmid:16002379 [CrossRef](#) [PubMed](#) [Google Scholar](#)
15. [↗](#)Bellinger DC. A strategy for comparing the contributions of environmental chemicals and other risk factors to neurodevelopment of children. *Environ Health Perspect*. 2012;**120**(4):501–507pmid:22182676 [PubMed](#) [Google Scholar](#)
16. [↗](#)Needleman HL, Riess JA, Tobin MJ, Biesecker GE, Greenhouse JB. Bone lead levels and delinquent behavior. *JAMA*. 1996;**275**(5):363–369pmid:8569015 [CrossRef](#) [PubMed](#) [Google Scholar](#)
17. [↗](#)Froehlich TE, Lanphear BP, Auinger P, et al. The association of tobacco and lead exposure with attention-deficit/hyperactivity disorder. *Pediatrics*. 2009;**124**(6). Available at: www.pediatrics.org/cgi/content/full/124/6/e1054pmid:19933729 [Abstract/FREE Full Text](#) [Google Scholar](#)
18. [↗](#)Nigg JT, Knottnerus GM, Martel MM, et al. Low blood lead levels associated with clinically diagnosed attention-deficit/hyperactivity disorder and mediated by weak cognitive control. *Biol Psychiatry*. 2008;**63**(3):325–331pmid:17868654 [CrossRef](#) [PubMed](#) [Google Scholar](#)
19. [↗](#)Dietrich KN, Ris MD, Succop PA, Berger OG, Bornschein RL. Early exposure to lead and juvenile delinquency. *Neurotoxicol Teratol*. 2001;**23**(6):511–518pmid:11792521 [CrossRef](#) [PubMed](#) [Google Scholar](#)
20. [↗](#)Wright JP, Dietrich KN, Ris MD, et al. Association of prenatal and childhood blood lead concentrations with criminal arrests in early adulthood. *PLoS Med*. 2008;**5**(5):e101pmid:18507497 [CrossRef](#) [PubMed](#) [Google Scholar](#)
21. [↗](#)Fergusson DM, Boden JM, Horwood LJ. Dentine lead levels in childhood and criminal behaviour in late adolescence and early adulthood. *J Epidemiol Community Health*. 2008;**62**(12):1045–1050pmid:18413435 [Abstract/FREE Full Text](#) [Google Scholar](#)
22. [↗](#)Marcus DK, Fulton JJ, Clarke EJ. Lead and conduct problems: a meta-analysis. *J Clin Child Adolesc Psychol*. 2010;**39**(2):234–241pmid:20390814 [CrossRef](#) [PubMed](#) [Google Scholar](#)
23. [↗](#)Reyes JW. Environmental policy as social policy? The impact of childhood lead exposure on crime. *BE J Econ Anal Policy*. 2007;**7**(1):1–41 [Google Scholar](#)
24. [↗](#)Fadrowski JJ, Navas-Acien A, Tellez-Plaza M, Guallar E, Weaver VM, Furth SL. Blood lead level and kidney function in US adolescents: the Third National Health and Nutrition Examination Survey. *Arch Intern Med*. 2010;**170**(1):75–82pmid:20065202 [CrossRef](#) [PubMed](#) [Google Scholar](#)
25. [↗](#)Borja-Aburto VH, Hertz-Picciotto I, Rojas Lopez M, Farias P, Rios C, Blanco J. Blood lead levels measured prospectively and risk of spontaneous abortion. *Am J Epidemiol*. 1999;**150**(6):590–597pmid:10489998 [Abstract/FREE Full Text](#) [Google Scholar](#)

26. ↵Zhu M, Fitzgerald EF, Gelberg KH, Lin S, Druschel CM. Maternal low-level lead exposure and fetal growth. *Environ Health Perspect*. 2010;**118**(10):1471–1475pmid:20562053 [CrossRef](#) [PubMed](#) [Google Scholar](#)
27. ↵Jones RL, Homa DM, Meyer PA, et al. Trends in blood lead levels and blood lead testing among US children aged 1 to 5 years, 1988–2004. *Pediatrics*. 2009;**123**(3). Available at: www.pediatrics.org/cgi/content/full/123/3/e376pmid:19254973 [Abstract/FREE Full Text](#) [Google Scholar](#)
28. ↵Dietrich KN, Ware JH, Salganik M, et al; Treatment of Lead-Exposed Children Clinical Trial Group. Effect of chelation therapy on the neuropsychological and behavioral development of lead-exposed children after school entry. *Pediatrics*. 2004;**114**(1):19–26pmid:15231903 [Abstract/FREE Full Text](#) [Google Scholar](#)
29. ↵Trasande L, Liu Y. Reducing the staggering costs of environmental disease in children, estimated at \$76.6 billion in 2008. *Health Aff (Millwood)*. 2011;**30**(5):863–870pmid:21543421 [Abstract/FREE Full Text](#) [Google Scholar](#)
30. ↵Gould E. Childhood lead poisoning: conservative estimates of the social and economic benefits of lead hazard control. *Environ Health Perspect*. 2009;**117**(7):1162–1167pmid:19654928 [CrossRef](#) [PubMed](#) [Google Scholar](#)
31. ↵Lanphear BP, Hornung R, Ho M, Howard CR, Eberly S, Knauf K. Environmental lead exposure during early childhood [published correction appears in *J Pediatr*. 2002;140(4):490]. *J Pediatr*. 2002;**140**(1):40–47pmid:11815762 [CrossRef](#) [PubMed](#) [Google Scholar](#)
32. ↵Ziegler EE, Edwards BB, Jensen RL, Mahaffey KR, Fomon SJ. Absorption and retention of lead by infants. *Pediatr Res*. 1978;**12**(1):29–34pmid:643372 [CrossRef](#) [PubMed](#) [Google Scholar](#)
33. ↵Wright RO, Shannon MW, Wright RJ, Hu H. Association between iron deficiency and low-level lead poisoning in an urban primary care clinic. *Am J Public Health*. 1999;**89**(7):1049–1053pmid:10394314 [CrossRef](#) [PubMed](#) [Google Scholar](#)
34. ↵US Department of Health and Human Services. *American Healthy Homes Survey. Lead and Arsenic Findings. Office of Healthy Homes and Lead Hazard Controls*. Washington, DC: US Department of Health and Human Services; 2011 [Google Scholar](#)
35. ↵Clark CS, Bornschein RL, Succop P, Que Hee SS, Hammond PB, Peace B. Condition and type of housing as an indicator of potential environmental lead exposure and pediatric blood lead levels. *Environ Res*. 1985;**38**(1):46–53pmid:4076111 [PubMed](#) [Google Scholar](#)
36. ↵US Department of Housing and Urban Development. *Guidelines for the Evaluation and Control of Lead-Based Paint Hazards in Housing*. 2nd ed. Washington, DC: US Department of Housing and Urban Development; 2012 [Google Scholar](#)
37. ↵US Environmental Protection Agency. 40 CFR part 745. Lead; identification of dangerous levels of lead: final rule. *Fed Regist*. 2001;**66**(4):1206–1240 [Google Scholar](#)
38. ↵Lanphear BP. The paradox of lead poisoning prevention. *Science*. 1998;**281**(5383):1617–1618pmid:9767027 [FREE Full Text](#) [Google Scholar](#)
39. Sayre JW, Charney E, Vostal J, Pless IB. House and hand dust as a potential source of childhood lead exposure. *Am J Dis Child*. 1974;**127**(2):167–170pmid:4810272 [CrossRef](#) [PubMed](#) [Google Scholar](#)
40. ↵Lanphear BP, Matte TD, Rogers J, et al. The contribution of lead-contaminated house dust and residential soil to children's blood lead levels. A pooled analysis of 12 epidemiologic studies. *Environ Res*. 1998;**79**(1):51–68pmid:9756680 [PubMed](#) [Google Scholar](#)

41. ↵ Aschengrau A, Beiser A, Bellinger D, Copenhafer D, Weitzman M. Residential lead-based-paint hazard remediation and soil lead abatement: their impact among children with mildly elevated blood lead levels. *Am J Public Health*. 1997;**87**(10):1698–1702pmid:9357358 [CrossRef](#) [PubMed](#) [Google Scholar](#)
42. ↵ Dixon SL, Gaitens JM, Jacobs DE, et al. Exposure of US children to residential dust lead, 1999–2004: II. The contribution of lead-contaminated dust to children's blood lead levels. *Environ Health Perspect*. 2009;**117**(3):468–474pmid:19337524 [CrossRef](#) [PubMed](#) [Google Scholar](#)
43. ↵ Amitai Y, Brown MJ, Graef JW, Cosgrove E. Residential deleading: effects on the blood lead levels of lead-poisoned children. *Pediatrics*. 1991;**88**(5):893–897pmid:1945628 [Abstract/FREE Full Text](#) [Google Scholar](#)
44. ↵ Clark S, Grote J, Wilson J, et al. Occurrence and determinants of increases in blood lead levels in children shortly after lead hazard control activities. *Environ Res*. 2004;**96**(2):196–205pmid:15325880 [PubMed](#) [Google Scholar](#)
45. ↵ Spanier AJ, Wilson S, Ho M, Hornung R, Lanphear BP. The contribution of housing renovation to children's blood lead levels: a cohort study. *Environ Health*. 2013;**12**:72pmid:23981571 [CrossRef](#) [PubMed](#) [Google Scholar](#)
46. ↵ Shannon M, Graef JW. Lead intoxication from lead-contaminated water used to reconstitute infant formula. *Clin Pediatr (Phila)*. 1989;**28**(8):380–382pmid:2758720 [Abstract/FREE Full Text](#) [Google Scholar](#)
47. ↵ Edwards M, Triantafyllidou S, Best D. Elevated blood lead in young children due to lead-contaminated drinking water: Washington, DC, 2001–2004. *Environ Sci Technol*. 2009;**43**(5):1618–1623pmid:19350944 [CrossRef](#) [PubMed](#) [Google Scholar](#)
48. ↵ Hanna-Attisha M, LaChance J, Sadler RC, Champney Schnepf A. Elevated blood lead levels in children associated with the Flint drinking water crisis: a spatial analysis of risk and public health response. *Am J Public Health*. 2016;**106**(2):283–290pmid:26691115 [CrossRef](#) [PubMed](#) [Google Scholar](#)
49. ↵ Richmond-Bryant J, Meng Q, Davis A, et al. The influence of declining air lead levels on blood lead–air lead slope factors in children. *Environ Health Perspect*. 2014;**122**(7):754–760pmid:24667492 [PubMed](#) [Google Scholar](#)
50. ↵ Levin R, Brown MJ, Kashtock ME, et al. Lead exposures in U.S. children, 2008: implications for prevention. *Environ Health Perspect*. 2008;**116**(10):1285–1293pmid:18941567 [CrossRef](#) [PubMed](#) [Google Scholar](#)
51. Centers for Disease Control and Prevention (CDC). Lead poisoning in pregnant women who used Ayurvedic medications from India: New York City, 2011–2012. *MMWR Morb Mortal Wkly Rep*. 2012;**61**(33):641–646pmid:22914225 [PubMed](#) [Google Scholar](#)
52. ↵ Gorospe EC, Gerstenberger SL. Atypical sources of childhood lead poisoning in the United States: a systematic review from 1966–2006. *Clin Toxicol (Phila)*. 2008;**46**(8):728–737pmid:18608287 [CrossRef](#) [PubMed](#) [Google Scholar](#)
53. ↵ Roscoe RJ, Gittleman JL, Deddens JA, Petersen MR, Halperin WE. Blood lead levels among children of lead-exposed workers: a meta-analysis. *Am J Ind Med*. 1999;**36**(4):475–481pmid:10470013 [CrossRef](#) [PubMed](#) [Google Scholar](#)
54. ↵ US Environmental Protection Agency. 40 CFR part 745. Lead; Clearance and clearance testing requirements for the renovation, repair and painting program. *Fed Regist*. 2010;**75**(87):25037–25073 [Google Scholar](#)
55. ↵ Farfel MR, Rohde C, Lees PSJ, Rooney B, Bannon DL, Derbyshire W. *Lead-Based Paint Abatement and Repair and Maintenance Study in Baltimore: Findings Based on Two Years of Follow-Up*. Washington, DC: US Environmental Protection Agency; 1998 [Google Scholar](#)

56. ↵ Galke W, Clark S, Wilson J, et al. Evaluation of the HUD lead hazard control grant program: early overall findings. *Environ Res*. 2001;**86**(2):149–156pmid:11437461 [PubMed](#) [Google Scholar](#)
57. ↵ Del Toral MA, Porter A, Schock MR. Detection and evaluation of elevated lead release from service lines: a field study. *Environ Sci Technol*. 2013;**47**(16):9300–9307pmid:23879429 [CrossRef](#) [PubMed](#) [Google Scholar](#)
58. ↵ Schock MR. Causes of temporal variability of lead in domestic plumbing systems. *Environ Monit Assess*. 1990;**15**(1):59–82pmid:24243429 [CrossRef](#) [PubMed](#) [Google Scholar](#)
59. Schock MR, Lemieux FG. Challenges in addressing variability of lead in domestic plumbing. *Water Science & Technology: Water Supply*. 2010;**10**(5):792–798 [Google Scholar](#)
60. Schock MR, Lytle DA. *Water Quality and Treatment: A Handbook of Community Water Supplies*. 6th ed. New York, NY: McGraw-Hill Inc; 2011 [Google Scholar](#)
61. ↵ Schock MR, Sandvig AM, Lemieux FG, DeSantis MK. Diagnostic sampling to reveal hidden lead and copper health risks. Presented at the *15th Canadian National Conference and 6th Policy Forum on Drinking Water*; Kelowna, BC; October 21–24, 2012 [Google Scholar](#)
62. ↵ Ossiander EM. A systematic review of screening questionnaires for childhood lead poisoning. *J Public Health Manag Pract*. 2013;**19**(1):E21–E29pmid:22668673 [CrossRef](#) [PubMed](#) [Google Scholar](#)
63. ↵ Centers for Disease Control. *Preventing Lead Poisoning in Young Children: A Statement by the Centers for Disease Control and Prevention*. Atlanta, GA: US Department of Health and Human Services; 1991 [Google Scholar](#)
64. ↵ Centers for Disease Control and Prevention. *Screening Young Children for Lead Poisoning: Guidance for State and Local Public Health Officials*. Atlanta, GA: Centers for Disease Control and Prevention; 1997 [Google Scholar](#)
65. ↵ Karp R, Abramson J, Clark-Golden M, et al. Should we screen for lead poisoning after 36 months of age? Experience in the inner city. *Ambul Pediatr*. 2001;**1**(5):256–258pmid:11888411 [CrossRef](#) [PubMed](#) [Google Scholar](#)
66. ↵ Clinical Laboratory Improvement Amendments of 1988. Pub L No. 100-578, 102 Stat 2903, 10 USC §263a (1988)
67. ↵ Parsons PJ, Geraghty C, Verostek MF. An assessment of contemporary atomic spectroscopic techniques for the determination of lead in blood and urine matrices. *Spect Act B*. 2001;**56**(9):1593–1604 [CrossRef](#) [Google Scholar](#)
68. ↵ Centers for Disease Control and Prevention. *Guidelines for the Identification and Management of Lead Exposure in Pregnant and Lactating Women*. Atlanta, GA: Centers for Disease Control and Prevention; 2010 [Google Scholar](#)
69. ↵ American College of Obstetricians and Gynecologists. Committee opinion no. 533. Lead screening during pregnancy and lactation. *Obstet Gynecol*. 2012;**120**(2 Pt 1):416–420pmid:22825110 [CrossRef](#) [PubMed](#) [Google Scholar](#)
70. ↵ Centers for Disease Control and Prevention. *Preventing Lead Exposure in Young Children: A Housing-Based Approach to Primary Prevention of Lead Poisoning*. Atlanta, GA: Centers for Disease Control and Prevention; 2004 [Google Scholar](#)
71. ↵ Geltman PL, Brown MJ, Cochran J. Lead poisoning among refugee children resettled in Massachusetts, 1995 to 1999. *Pediatrics*. 2001;**108**(1):158–162pmid:11433069 [Abstract/FREE Full Text](#) [Google Scholar](#)
72. ↵ Centers for Disease Control and Prevention (CDC). Elevated blood lead levels among internationally adopted children: United States, 1998. *MMWR Morb Mortal Wkly Rep*. 2000;**49**(5):97–100pmid:10718094 [PubMed](#) [Google Scholar](#)

[Advertising Disclaimer »](#)

In this issue

Pediatrics

Vol. 138, Issue 1

1 Jul 2016

[Table of Contents](#)

[Index by author](#)

View this article with *LENS*

[✉ Email Article](#)

[↻ Share](#)

[© Request Permissions](#)

[🖨 Print](#)

[! Article Alerts](#)

[📄 Download PDF](#)

[🔍 Citation Tools](#)

[🔔 Insight Alerts](#)

[Table of Contents](#)

Related Articles

Boost efforts to reduce lead poisoning before children are exposed: AAP policy

Bruce P. Lanphear et al., AAP News, 2016

Still Treating Lead Poisoning After All These Years

Bruce Lanphear, Pediatrics, 2017

Lead Intoxication in Infancy

Michael W. Shannon et al., Pediatrics, 1992

Survey of Lead Exposure Around a Closed Lead Smelter

Renate Kimbrough et al., Pediatrics, 1995

Lead Exposure in Children: Prevention, Detection, and Management

Committee on Environmental Health, Pediatrics, 2005

The Paradox of Lead Poisoning Prevention

Bruce P. Lanphear, Science, 1998

Environmental racism and the need for private well protections

Anne E. Nigra, Proc Natl Acad Sci U S A, 2020

No Benefit From Lowering Lead

Science, 2001

Measuring lead health risks

Pigment & Resin Technology, 1998

Lead in paint and dust from a children's nursery

Jonathan M. Horner, Environmental Management and Health, 1995

Powered by **TREND MD**

▼ **Related Articles**

COUNCIL ON ENVIRONMENTAL HEALTH. Prevention of Childhood Lead Toxicity. *Pediatrics*. 2016;38(1):e20161493

Prevention of Childhood Lead Toxicity

AAP Publications Reaffirmed or Retired

[PubMed](#) [Google Scholar](#)

► **Cited By...**

► **More in this TOC Section**

► **Similar Articles**

Subjects

Environmental Health ▶

Environmental Health ▶

Lead ▶

Current Policy ▶

AAP Policy Collections by Authoring Entities ▶

Council on Environmental Health ▶



[See more details](#)

-  Picked up by **66** news outlets
-  Blogged by **6**
-  Referenced in **5** policy sources
-  Tweeted by **54**
-  On **7** Facebook pages
-  Mentioned in **1** Google+ posts
-  **230** readers on Mendeley

Journal Info

[Editorial Board](#)

[Editorial Policies](#)

[Overview](#)

[Licensing Information](#)

Authors/Reviewers

[Author Guidelines](#)

[Submit My Manuscript](#)

[Open Access](#)

[Reviewer Guidelines](#)

Librarians

[Institutional Subscriptions](#)

[Usage Stats](#)

Support

[Contact Us](#)

[Subscribe](#)

Resources

[Media Kit](#)

About

[International Access](#)

[Terms of Use](#)

[Privacy Statement](#)

[FAQ](#)

[AAP.org](#)

[shopAAP](#)



© 2021 American Academy of Pediatrics